



190 Campus Blvd, Ste. 300 • Winchester, VA 22601
(540)667-1244 • www.winchestergastro.com

**PREP INSTRUCTIONS for Upper Endoscopy,
Dilation, PEG or ERCP**

Arrive at the Endoscopy Center at the **Winchester Medical Center, Medical Office Building II, Ste. 100, Mall Level** at _____ on _____ for your procedure at _____. The Endoscopy center is located on the first floor of MOB II Ste. 100, Phone (540)536-8746. Do not go to the office on the 3rd floor. You are to arrive one hour in advance of the scheduled procedure time to allow for administrative check in and final procedure preparation. Procedure times are approximate and even though we intend to start on time there can be delays and we ask for your patience. On average you will be in the unit approximately 2-3 hours. Most procedures take 15 minutes and recovery from the sedation takes 45-60 minutes. Written discharge instructions will be given to you. You should take it easy after discharge until all the drug effects have worn off. Any diet instructions will be discussed with you by your physician prior to discharge. Any other special needs (like work slips), should be discussed prior to discharge with the nurse or physician. You will have a private holding room during your procedure which will accommodate comfortably one other individual, preferably your designated driver. You will not be allowed to drive yourself home because of the sedation used.

Diet

Your diet should be restricted for this procedure to allow for optimal visualization of your gastrointestinal tract. You may drink clear liquids in limited quantities up to 4 hours before the procedure. **NO SOLIDS DAY OF PROCEDURE. DAY OF PROCEDURE Clear liquids only** – no fatty liquids (no cream, no milk, no solids)

- DO NOT SMOKE, VAPE OR USE OTHER TOBACCO PRODUCTS THE DAY OF YOUR PROCEDURE

Medications

You may take your usual medications with a small amount of water no later than 2 hours before the scheduled procedure time. Diabetes medications are usually held or reduced to half strength before these procedures because your food intake is being restricted. This should be discussed at the time the procedure is scheduled. Your medications can usually be restarted immediately after the procedure.

*If taking Phentermine stop taking 7 days prior to your procedure.

Anticoagulants

Aspirin, Plavix, and Coumadin should not be stopped for routine upper endoscopy. They are usually stopped for esophageal dilations and variceal banding. Patients having these procedures should discuss how to handle the blood thinners at the time the procedure is scheduled. Patients who cannot stop blood thinners because of the high risk of stroke should discuss any changes of blood thinners with your doctor.

Antibiotics are no longer required for routine endoscopic procedures. PEG and ERCP are the exceptions.

****IF YOU CAN NOT MAKE YOUR APPOINTMENT TIME, PLEASE IMMEDIATELY NOTIFY US AT 540 667-1244. THIS NUMBER IS AVAILABLE AFTERHOURS FOR MESSAGES. FAILURE TO NOTIFY US FOR CANCELLATION WITHIN 48 HOURS MAY RESULT IN A CHARGE OF \$75.00 TO YOU. ****

*****INSURANCE: WE WILL CONTACT YOUR INSURANCE CARRIER FOR PRE-AUTHORIZATION; HOWEVER, IT IS YOUR RESPONSIBILITY TO CHECK ON THE BENEFITS FOR COVERED SERVICES FROM YOUR INSURANCE CARRIER. WE MAY USE AN ANESTHESIOLOGIST FOR SEDATION, WHICH COULD INCURE AN ADDITIONAL COST/BILL. PLEASE CONTACT OUR**

Llewellyn Kitchin, MD • Richard Sears, MD • Richard Nemec, MD • Grace Shih, MD • Khurram Baig, DO •
Satyanisth Agrawal, DO

**OFFICE IF YOU CHOOSE NOT TO USE AN ANESTHESIOLOGIST FOR SEDATION. IF YOU DO NOT HAVE INSURANCE, PLEASE CALL OUR OFFICE AT (540)
667-1244 TO MAKE PAYMENT ARRANGEMENTS. *****